

## Effects of neurodynamic mobilization versus foam rolling treatment after delayed-onset muscle soreness



B. Romero-Moraleda<sup>a,\*,</sup> D. Muñoz-García<sup>b,</sup> S. Lerma Lara<sup>c,</sup> R. Ferrer Peña<sup>d,</sup> V. Paredes<sup>e,</sup> A.B. Peinado<sup>f</sup>

<sup>a</sup> CSEU La Salle-UAM, Health Sciences Faculty, Camilo José Cela University, Madrid, Spain

<sup>b</sup> CSEU La Salle-UAM, Group on Movement-Behavioral Science and Study of Pain, Madrid, Spain

<sup>c</sup> CSEU La Salle-UAM, Hospital Infantil Universitario Niño Jesús, Madrid, Spain

<sup>d</sup> CSEU La Salle-UAM, Centro de Atención Primaria, Entrevías, Madrid, Spain

<sup>e</sup> Health Sciences Faculty, Camilo José Cela University, Rayo Vallecano S.A.D., Villanueva de la Cañada, Madrid, Spain

<sup>f</sup> School of Physical Activity and Sport Sciences, LFE Research Group, Technical University of Madrid, Madrid, Spain

**Research question:** Is neurodynamic mobilization better than a foam rolling treatment for treating muscle induced pain?

**Introduction:** Delayed-onset muscle soreness (DOMS) refers to the skeletal muscle pain that is experienced following eccentric exercise [1]. Exercise-induced muscle damage is known to be manifested as increased serum concentrations of creatine kinase (CK), ultrastructural disruption, inflammation and strength deficits [2]. Numerous recovery strategies have been used in an attempt to minimize the symptoms of DOMS [3]. The purpose of this study was to assess the acute effects of a single session of NM treatment and to compare with foam roller application after exercise-induced muscle damage [4].

**Materials and methods:** Thirty-two healthy subjects (21 males and 11 females, mean age:  $22.6 \pm 2.2$  years) were randomly assigned into neurodynamic group (ND,  $n = 16$ ) and foam roller group (FR,  $n = 16$ ). Drop jumps were used to induce muscle damage (5 set  $\times$  20 rep/2 min. recovery). Maximum voluntary contraction (MIVC), muscle activation in landing jump (MALJ), 0–10 pain scale, pressure pain thresholds, hip and knee ROM and knee bend neural test were measured baseline, post 48 h pre-treatment and immediately post treatment. A PhysioPlux<sup>®</sup> system was used for sEMG data collection. Electrode placement was done following the SENIAM recommendations. A band-pass filter was performed to the raw signals and RMS was calculated. A digital algometer (Wagner Pain Test<sup>®</sup>) was used for PPT and an universal goniometer for ROM measurements.

**Results:** MIVC, MALJ and hip and knee ROM were decreased and DOMS measured through pain increased in both groups ( $p < 0.05$ ). After treatment, both groups showed improvements for strength and pain. Pain scale decreased significantly without differences between groups (Pain FR:  $-2.38 \pm 1.41\%$ ; Pain ND:  $-2.53 \pm 0.92\%$ ,  $p = 0.71$ ). For strength measures, the FR achieved greatest changes than ND (MIVC FR:  $10.8 \pm 4.65\%$ ; MIVC ND:  $4.80 \pm 4.33\%$ ;  $p = 0.03$ ).

**Discussion:** The data indicated that the both treatments were effective to improve pain and strength, showing EMG measures greater change with FR treatment.

### References

- [1] Kanda K, Sugama K, Hayashida H, Sakuma J, Kawakami Y, Miura S, et al. Eccentric exercise-induced delayed-onset muscle soreness and changes in markers of muscle damage and inflammation. *Exerc Immunol Rev* 2013;19:72–85.
- [2] Miyama M, Nosaka K. Influence of surface on muscle damage and soreness induced by consecutive drop jumps. *J Strength Cond Res* 2004;18(2):206–11.
- [3] Torres R, Ribeiro F, Akbeto Duarte J, Cabri JM. Evidence of the physiotherapeutic interventions used currently after exercise-induced muscle damage: systematic review and meta-analysis. *Phys Ther Sport* 2012;13(2):101–14.
- [4] Howatson G, Goodall S, Van Someren K. The influence of cold water immersions on adaptation following a single bout of damaging exercise. *Eur J Appl Physiol* 2009;105(4):615–21.